



ADAMA

MAXENTIS®

Provides protection against white mould in soybeans and features two distinct, elite modes of action for resistance management (Groups 3 & 11) offering long-lasting, systemic activity.



POWERED BY

Asorbital

FORMULATION TECHNOLOGY

Active Ingredients

Azoxystrobin 120 g/L +
Prothioconazole 90 g/L
as an EC

Packaging

Case: 2 x 8.45 L jugs

Application Rates and Acres Treated

Rate: 422 ml/ac

Acres Treated: 20 ac/jug

Water Volume

Ground: 40 L/ac (10 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

Rainfastness

Avoid applying when rain is forecast.

GROUPS 3 & 11

KEY BENEFITS

- Unique combination of prothioconazole and azoxystrobin
- Enhanced EC formulation – with bulk packaging capabilities
- Multi-mode activity for resistance management (Groups 3 & 11)
- Broad-spectrum disease control with protective and curative action
- Whole plant protection: translaminar and systemic movement
- Wide window of application

REGISTERED CROPS*

- Soybeans

APPLICATION TIMING AND CROP STAGING (SOYBEANS)

Diseases*	Rate (ml/ac)	Timing
White mould (<i>Sclerotinia</i>)	422	Apply as a preventive foliar spray at the beginning of flowering or at first sign of disease. After the initial application, one additional application may be made 10–14 days afterwards if conditions remain favourable for continued or increased disease development. Apply the higher rate when conditions favour disease development, or when growing less disease resistant varieties. Maximum applications per year: 2

* See label for the complete list of registered crops as well diseases controlled and rates for each crop.



MAXENTIS®

HOW IT WORKS

Two powerful active ingredients, combined for the first time in Canada, that provide protective and curative action on a wide range of diseases, across multiple crops.

MIXING INSTRUCTIONS

1. Fill the clean spray tank $\frac{3}{4}$ full of clean water.
2. Add the required amount of MAXENTIS® and agitate thoroughly.
3. Continue agitation and add the required amount of the tank-mix partner.
4. Complete filling tank to the desired level with water.


CROP ROTATIONS

No restrictions

TRIAL DATA

2023 Soybean Yield Ontario (bu/ac)

2 SITES



QUICK TIP:

Environmental conditions are a major driver of disease pressure and severity. Under high disease pressure conditions a rate of MAXENTIS® at the upper end of the rate range is recommended.

GRAZING RESTRICTIONS

Forage, hay: 30 days

Grazing or green feed: 6 days

STORAGE

Do not freeze.


PRE-HARVEST INTERVAL (SOYBEANS)

20 days

NOTE: Refer to the label for pre-harvest intervals for all other registered crops.

WANT TO KNOW MORE?

Click this QR for full product information including label and SDS documents.



ADAMA

1.855.264.6262 ADAMA.COM

®/™ ASORBITAL® and MAXENTIS® are registered trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.

© 2023 ADAMA Agricultural Solutions Canada Ltd.

05/2024