# **MAXENTIS®**

ADAMA

Provides protection against the most damaging diseases in peas, lentils and canola. It features two distinct, elite modes of action for resistance management (Groups 3 & 11) along with long-lasting systemic activity.

#### POWERED BY Asorbital® FORMULATION TECHNOLOGY

#### **Active Ingredients:**

Azoxystrobin 120 g/L + Prothioconazole 90 g/L as an EC

#### **Packaging**:

MAXENTI

Case: 2 x 8.45 L jugs Drum: 118.1 L

#### **Application Rates** and Acres Treated:

CANOLA: Rate: 253 – 443 ml/ac Acres Treated: 19 – 33 ac/jug; 267 – 467 ac/drum

#### PEAS, LENTILS & SOYBEANS:

Rate: 422ml/ac Acres Treated: 20 ac/jug; 280 ac/drum

#### Water Volume:

Ground: 40 L/ac (10 US gal/ac) Aerial: 20 L/ac (5 US gal/ac)

#### **Rainfastness:**

Avoid applying when rain is forecast.

#### GROUPS 3 & 11

#### **KEY BENEFITS:**

- Unique combination of prothioconazole and azoxystrobin
- Enhanced EC formulation with bulk packaging capabilities
- Multi-mode activity for resistance management (Groups 3 & 11)

#### **REGISTERED CROPS\*:**

- Canola
- Peas

- Broad-spectrum disease control with protective and curative action
- Whole plant protection: translaminar and systemic movement
- Wide window of application
- Lentils
- Soybeans

#### **KEY DISEASES CONTROLLED (OILSEEDS AND PULSES)\*:**

- Anthracnose
- Ascochyta Blight
- Blackleg

- Mycosphaerella Blight (suppression)
- White Mould (Sclerotinia)

#### **HOW IT WORKS:**

Two powerful active ingredients, combined for the first time in Canada, that provide protective and curative action on a wide range of diseases, across multiple crops.

\* See label for the complete list of registered crops as well diseases controlled and rates for each crop.

## **MAXENTIS**®

#### APPLICATION TIMING AND CROP STAGING (PULSES AND OILSEEDS):

Сгор	Diseases	Rate (ml/ac)	Timing
Canola	White Mould (Sclerotinia)	443	Apply when the crop is in the 20–50% bloom stage. Best protection will be achieved when the fungicide is applied prior to petals beginning to fall and will allow for the maximum number of petals to be protected. Maximum applications per year: 1 Blackleg: Early application required 2-6 leaf
	Blackleg	253	
Peas	Mycosphaerella Blight (suppression) White Mould (Sclerotinia)	422	Apply as a preventive foliar spray at the beginning of flowering or at first sign of disease. After the initial application, one additional application may be made 10–14 days afterwards if conditions remain favourable for continued or increased disease development. Apply the higher rate when conditions favour disease development, or when growing less disease resistant varieties.
Lentils	Anthracnose, Ascochyta Blight, White Mould		
Soybeans	White Mould (Sclerotinia)		Maximum applications per year: 2

#### **MIXING INSTRUCTIONS:**

- 1. Fill the clean spray tank ¾ full of clean water.
- 2. Add the required amount of MAXENTIS® and agitate thoroughly.
- Continue agitation and add the required amount of the tank-mix partner.
- 4. Complete filling tank to the desired level with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

#### **GRAZING RESTRICTIONS:**

Forage, Hay: 30 days Grazing or Green Feed: 6 days

NOTE: Do not feed dried pea vines to livestock.

#### **CROP ROTATIONS:**

No restrictions

#### **PRE-HARVEST INTERVALS:**

- Peas, Lentils: 15 days
- Soybeans: 20 days
- Canola: 36 days

NOTE: Refer to the label for pre-harvest intervals for all other registered crops.

#### **STORAGE:**

Do not freeze.



### **QUICK TIPS:**

Environmental conditions are a major driver of disease pressure and severity. Under high disease pressure conditions a rate of MAXENTIS® at the upper end of the rate range is recommended.

ADAMA

Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. 1.855.264.6262 | ADAMA.COM

MAXENTIS® is a registered trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.